arXiv:1410.2668v1 [math.NT] 10 Oct 2014

Images of 2-adic representations associated to hyperelliptic Jacobians

Jeffrey Yelton

October 13, 2014

Abstract

Let k be a subfield of $\mathbb{C}$ which contains all 2-power roots of unity, and let $K = k(\alpha_1, \alpha_2, ..., \alpha_{2g+1})$, where the α_i's are independent and transcendental over k, and g is a positive integer. We investigate the image of the 2-adic Galois action associated to the Jacobian J of the hyperelliptic curve over K given by $y^2 = \prod_{i=1}^{2g+1}(x - \alpha_i)$. Our main result states that the image of Galois in $\mathrm{Sp}(T_2(J))$ coincides with the principal congruence subgroup $\Gamma(2) \lhd \mathrm{Sp}(T_2(J))$. As an application, we find generators for the algebraic extension $K(J[4])/K$ generated by coordinates of the 4-torsion points of J.

1 Introduction

Fix a positive integer g. An affine model for a hyperelliptic curve over $\mathbb{C}$ of genus g may be given by

$$y^2 = \prod_{i=1}^{2g+1}(x - \alpha_i), \tag{1}$$

with α_i's distinct complex numbers. Now let $\alpha_1, ..., \alpha_{2g+1}$ be transcendental and independent over $\mathbb{C}$, and let L be the subfield of $\mathbb{C}(\alpha) := \mathbb{C}(\alpha_1, ..., \alpha_{2g+1})$ generated over $\mathbb{C}$ by the elementary symmetric functions of the α_i's. For any positive integer N, let $J[N]$ denote the N-torsion subgroup of $J(\bar{L})$. For each $n \geq 0$, let $L_n = L(J[2^n])$ denote the extension of L over which the 2^n-torsion of J is defined. Set

$$L_\infty := \bigcup_{n=1}^{\infty} L_n.$$

Note that $\mathbb{C}(\alpha_1, ...\alpha_{2g+1})$ is Galois over L with Galois group isomorphic to S_{2g+1}. It is well known ([5], Corollary 2.11) that $\mathbb{C}(\alpha_1, ..., \alpha_{2g+1}) = L_1$, so $\mathrm{Gal}(L_1/L) \cong S_{2g+1}$. Fix an algebraic closure $\bar{L}$ of L, and write G_L for the absolute Galois group $\mathrm{Gal}(\bar{L}/L)$.

Let C be the curve defined over L by equation (1), and let J/L be its Jacobian. For any prime ℓ, let

$$T_\ell(J) := \lim_{\leftarrow n} J[\ell^n]$$

denote the ℓ-adic Tate module of J; it is a free $\mathbb{Z}_\ell$-module of rank $2g$ (see [6], §18). For the rest of this paper, we write $\rho_\ell : G_L \to \mathrm{Aut}(T_\ell(J))$ for the continuous homomorphism induced by the natural Galois action on $T_\ell(J)$. Write $\mathrm{SL}(T_\ell(J))$ (resp. $\mathrm{Sp}(T_\ell(J))$) for the subgroup of automorphisms of the 2-adic Tate module $T_\ell(J)$ with determinant 1 (resp. automorphisms of $T_\ell(J)$ which preserve the Weil pairing). Since L contains all 2-power roots of unity, the Weil pairing on $T_2(J)$ is Galois invariant, and it follows that the image of ρ_2 is contained in $\mathrm{Sp}(T_2(J))$. For each $n \geq 0$, we denote by

$$\Gamma(2^n) := \{g \in \mathrm{Sp}(T_2(J)) \mid g \equiv 1 \pmod{2^n}\} \lhd \mathrm{Sp}(T_2(J))$$

the level-2^n principal congruence subgroup of $\mathrm{Sp}(T_2(J))$.

Our main theorem is the following.

Theorem 1.1. *With the above notation, the image under ρ_2 of the Galois subgroup fixing L_1 is $\Gamma(2) \lhd \mathrm{Sp}(T_2(J))$.*

Before setting out to prove this theorem, we state some easy corollaries.

Corollary 1.2. *Let G denote the image under ρ_2 of all of G_L. Then we have the following:*

a) G contains $\Gamma(2) \lhd \mathrm{Sp}(T_2(J))$, and $G/\Gamma(2) \cong S_{2g+1}$.

b) In the case that $g = 1$, $G = \mathrm{Sp}(T_2(J)) = \mathrm{SL}(T_2(J))$.

c) For each $n \geq 1$, the homomorphism ρ_2 induces an isomorphism

$$\bar{\rho}_2^{(n)} : \mathrm{Gal}(L_n/L_1) \xrightarrow{\sim} \Gamma(2)/\Gamma(2^n)$$

via the restriction map $\mathrm{Gal}(\bar{L}/L_1) \twoheadrightarrow \mathrm{Gal}(L_n/L_1)$.

Proof. Since $\mathrm{Gal}(L_1/L) \cong S_{2g+1}$, part (a) immediately follows from the theorem. If $g = 1$, then fix a basis of $T_2(J)$ so that we may identify $\mathrm{Sp}(T_2(J))$ (resp. $\mathrm{SL}(T_2(J))$) with $\mathrm{Sp}_2(\mathbb{Z}_2)$ (resp. $\mathrm{SL}_2(\mathbb{Z}_2)$). Then it is well known that $\mathrm{Sp}_2(\mathbb{Z}_2) = \mathrm{SL}_2(\mathbb{Z}_2)$, and that $\mathrm{SL}_2(\mathbb{Z}_2)/\Gamma(2) \cong \mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z}) \cong S_3$. Since, by part (a), $G/\Gamma(2) \cong S_3$ when $g = 1$, the linear subgroup G must be all of $\mathrm{Sp}(T_2(J)) = \mathrm{SL}(T_2(J))$, which is the statement of (b). To prove part (c), note that for any $n \geq 0$, the image under ρ_2 of the Galois subgroup fixing the 2^n-torsion points is clearly $G \cap \Gamma(2^n)$. But $G > \Gamma(2)$, so for any $n \geq 1$, the image under ρ_2 of $\mathrm{Gal}(\bar{L}/L(2^n))$ is $\Gamma(2^n)$. Then part (c) immediately follows by the definition of $\bar{\rho}_2^{(n)}$.

□

In §2, we will prove the main theorem by considering a family of hyperelliptic curves whose generic fiber is C. In §3, we will use the results of the previous two sections to determine generators for the algebraic extension L_2/L (Theorem 3.1). Finally, in §4, we will generalize Theorems 1.1 and 3.1 by descending from $\mathbb{C}$ to a subfield $k \subset \mathbb{C}$ which contains all 2-power roots of unity.

2 Families of hyperelliptic Jacobians

In order to prove Theorem 1.1, we study a family of hyperelliptic curves parametrized by all (unordered) $(2g+1)$-element subsets $T = \{\alpha_i\} \subset \mathbb{C}$ whose generic fiber is C. Let $e_1 := \sum_{i=1}^{2g+1} \alpha_i, ..., e_{2g+1} := \prod_{i=1}^{2g+1} \alpha_i$ be the elementary symmetric functions of the variables α_i, and let Δ be the discriminant function of these variables. Then the base of this family is the affine variety over $\mathbb{C}$ given by

$$X := \mathrm{Spec}(\mathbb{C}[e_1, e_2, ..., e_{2g+1}, \Delta^{-1}]). \tag{2}$$

This complex affine scheme may be viewed as the configuration space of $(2g+1)$-element subsets of $\mathbb{C}$ (see the discussion in Section 6 of [10]). More precisely, we identify each $\mathbb{C}$-point $T = (e_1, e_2, ..., e_{2g+1})$ of X with the set of roots of the squarefree degree-$(2g+1)$ polynomial $z^{2g+1} - e_1 z^{2g} + e_2 z^{2g-1} - ... - e_{2g+1} \in \mathbb{C}[z]$, which is a $(2g+1)$-element subset of $\mathbb{C}$. Note that the function field of X is L. The (topological) fundamental group of X is isomorphic to B_{2g+1}, the braid group on $2g+1$ strands. The braid group B_{2g+1} is generated by elements σ_1, σ_2, ... , σ_{2g}, with relations $\sigma_1\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\sigma_{i+1}$ for $1 \leq i \leq 2g$ and $\sigma_i\sigma_j = \sigma_j\sigma_i$ for $2 \leq i+1 < j \leq 2g$. (See section 1.4 of [2] for more details.)

We also define the complex affine scheme

$$Y := \mathrm{Spec}(\mathbb{C}[\alpha_1, \alpha_2, ..., \alpha_{2g+1}, \{(\alpha_i - \alpha_j)^{-1}\}_{1 \leq i < j \leq 2g+1}]). \tag{3}$$

As a complex manifold, Y is the *ordered* configuration space, whose $\mathbb{C}$-points may be identified with $2g+1$-element subsets of $\mathbb{C}$ which are given an ordering (a $\mathbb{C}$-point is identified with its coordinates $(\alpha_1, \alpha_2, ..., \alpha_{2g+1})$). There is an obvious covering map $Y \to X$ which sends each point $(\alpha_1, \alpha_2, ..., \alpha_{2g+1})$ of Y to the point in X corresponding to the (unordered) subset $\{\alpha_1, \alpha_2, ..., \alpha_{2g+1}\}$. The *pure* braid group on $2g+1$ strands, denoted P_{2g+1}, is defined to be the kernel of the surjective homomorphism from B_{2g+1} to the symmetric group S_{2g+1} which sends σ_i to $(i, i+1) \in S_{2g+1}$ for $1 \leq i \leq 2g$ (see the proof of Theorem 1.8 in [2]). Then $P_{2g+1} \lhd B_{2g+1}$ is the (normal) subgroup corresponding to the cover $Y \to X$, and is therefore isomorphic to the fundamental group of Y.

Let $\mathcal{O}_X$ denote the coordinate ring of X, and let $F(x) \in \mathcal{O}_X[x]$ be the degree-$(2g+1)$ polynomial given by

$$x^{2g+1} + \sum_{i=1}^{2g+1} (-1)^i e_i x^{2g+1-i}. \tag{4}$$

Now denote by $\mathcal{C} \to X$ the affine scheme defined by the equation $y^2 = F(x)$. Clearly, $\mathcal{C}$ is the family over X whose fiber over a point $T \in X(\mathbb{C})$ is the smooth affine hyperelliptic curve defined by $y^2 = \prod_{z \in T}(x - z)$, and the generic fiber of $\mathcal{C}$ is C/L. Fix a basepoint T_0 of X, and a basepoint P_0 of $\mathcal{C}_{T_0}$. Then we have a short exact sequence of fundamental groups

$$1 \to \pi_1(\mathcal{C}_{T_0}, P_0) \to \pi_1(\mathcal{C}, P_0) \to \pi_1(X, T_0) \to 1. \tag{5}$$

We now construct a continuous section $s : X \to \mathcal{C}$, following the proof of Lemma 6.1 and the discussion in [10], §6. For $i = 1, 2$, let $\mathcal{E}_i \to X$ be the affine scheme given by $\mathrm{Spec}(\mathcal{O}_X[x, y]/(y^i - F(x))[F(x)^{-1}])$. Then $\mathcal{E}_1 \to X$ is clearly the family of complex topological spaces whose fiber over a point $T \in X$ can be identified with $\mathbb{C} \setminus T$, and there is an obvious degree-2 cover $\mathcal{E}_2 \to \mathcal{E}_1$. Let $t : X \to \mathcal{E}_1$ be the continuous map of complex topological spaces which sends a point $T \in X$ to $\max_{z \in T}\{|z|\} + 1 \in \mathbb{C} \setminus T = \mathcal{E}_{1,T}$. This section then lifts to a section $\tilde{t} : X \to \mathcal{E}_2$. Define $s : X \to \mathcal{C}$ to be the composition of $\tilde{t}$ with the obvious inclusion map $\mathcal{E}_2 \hookrightarrow \mathcal{C}$. It is easy to check from the construction of s that it is a section of the family $\mathcal{C} \to X$.

The section s induces a monodromy action of $\pi_1(X, T_0)$ on $\pi_1(\mathcal{C}_{T_0}, P_0)$, which is given by $\sigma \in \pi_1(X)$ acting as conjugation by $s(\sigma)$ on $\pi_1(\mathcal{C}_{T_0}, P_0) \lhd \pi_1(\mathcal{C}, P_0)$. This induces an action of B_{2g+1} on the abelianization of $\pi_1(\mathcal{C}_{T_0}, P_0)$, the homology group $H_1(\mathcal{C}_{T_0}, \mathbb{Z})$, which is isomorphic to $\mathbb{Z}^{2g}$. We denote this action by

$$R : B_{2g+1} \cong \pi_1(X, T_0) \to \mathrm{Aut}(H_1(\mathcal{C}_{T_0}, \mathbb{Z})). \tag{6}$$

This action respects the intersection pairing on $\mathcal{C}_{T_0}$, so the image of R is actually contained in the corresponding subgroup of symplectic automorphisms $\mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z}))$.

The following theorem is proven in [1] (Théoréme 1), as well as in [5] (Lemma 8.12).

Theorem 2.1. *In the representation $R : B_{2g+1} \to \mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z}))$, the image of P_{2g+1} coincides with $\Gamma(2)$.*

Let $\widehat{B}_{2g+1}$ denote the profinite completion of $B_{2g+1} \cong \pi_1(X, T_0)$. Since X may be viewed as a scheme over the complex numbers, Riemann's Existence Theorem yields an isomorphism between its étale fundamental group $\pi_1^{\acute{e}t}(X, T_0)$ and $\widehat{B}_{2g+1}$ ([3], Exposé XII, Corollaire 5.2). Meanwhile, $\pi_1^{\acute{e}t}(X, T_0)$ is isomorphic to the Galois group $\mathrm{Gal}(L^{\mathrm{unr}}/L)$, where L^{unr} is the maximal extension of L unramified at all points of X. The representation $R : B_{2g+1} \to \mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z}))$ induces a homomorphism of profinite groups

$$R : \mathrm{Gal}(L^{\mathrm{unr}}/L) = \widehat{B}_{2g+1} \to \mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z}) \otimes \mathbb{Z}_\ell) \tag{7}$$

for any prime ℓ. Composing this map with the restriction homomorphism $G_L := \mathrm{Gal}(\bar{L}/L) \twoheadrightarrow \mathrm{Gal}(L^{\mathrm{unr}}/L)$ yields a map which we denote $R_\ell : G_L \to \mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z}) \otimes \mathbb{Z}_\ell)$. The following proposition will allow us to convert the above topological result into the arithmetic statement of Theorem 1.1.

Proposition 2.2. *Assume the above notation, and let ℓ be any prime. Then there is an isomorphism of $\mathbb{Z}_\ell$-modules $T_\ell(J) \xrightarrow{\sim} H_1(\mathcal{C}_{T_0}, \mathbb{Z}) \otimes \mathbb{Z}_\ell$ making the representations ρ_ℓ and R_ℓ isomorphic.*

Proof. We proceed in five steps.

Step 1: We switch from the affine curve C to a smooth compactification of C, which is defined as follows. Let C' be the (smooth) curve defined over L by

the equation

$$y'^2 = x' \prod_{i=1}^{2g+1} (1 - \alpha_i x'). \tag{8}$$

We glue the open subset of C defined by $x \neq 0$ to the open subset of C' defined by $x' \neq 0$ via the mapping

$$x' \mapsto \frac{1}{x},\ y' \mapsto \frac{y}{x^{g+1}},$$

and denote the resulting smooth, projective scheme by $\bar{C}$. (See [5], §1 for more details of this construction.) Let $\infty \in \bar{C}(L)$ denote the "point at infinity" given by $(x', y') = (0, 0) \in C'$. The curve $\bar{C}$ has smooth reduction over every point $T \in X$ and therefore can be extended in an obvious way to a family $\bar{\mathcal{C}} \to X$ whose generic fiber is $\bar{C}/L$. Note that $\bar{\mathcal{C}}_T$ is a smooth compactification of $\mathcal{C}_T$ for each $T \in X$. There is a surjective map $\pi_1(\mathcal{C}_{T_0}, P_0) \twoheadrightarrow \pi_1(\bar{\mathcal{C}}_{T_0}, \infty_{T_0})$ induced by the inclusion $\mathcal{C} \hookrightarrow \bar{\mathcal{C}}$. Note also that the section $s : X \to \mathcal{C} \subset \bar{\mathcal{C}}$ can be continuously deformed to the "constant section" $\bar{s} : X \to \bar{\mathcal{C}}$ sending each $T \in X$ to the point at infinity $\infty_T \in \mathcal{C}_T$. Therefore, $\bar{s}_* : \pi_1(X, T_0) \to \pi_1(\bar{\mathcal{C}}_{T_0}, \infty_{T_0})$ is the composition of s_* with the map $\pi_1(\mathcal{C}_T) \twoheadrightarrow \pi_1(\bar{\mathcal{C}}_T)$. In this way, we may view the action of $\pi_1(X, T_0)$ on $\pi_1(\mathcal{C}_{T_0}, P_0)^{\mathrm{ab}} = \pi_1(\bar{\mathcal{C}}_{T_0}, \infty_{T_0})^{\mathrm{ab}}$ as being induced by $\bar{s}_*$.

Step 2: We switch from (topological) fundamental groups to étale fundamental groups. Since X and $\mathcal{C}$, as well as $\mathcal{C}_T$ for each $T \in X$, can be viewed as a scheme over the complex numbers, Riemann's Existence Theorem implies that the étale fundamental groups of X, $\mathcal{C}$, and each $\mathcal{C}_T$ (defined using a choice of geometric base point $\bar{T}_0$ over T_0) are isomorphic to the profinite completions of their respective topological fundamental groups. Taking profinite completions induces a sequence of étale fundamental groups

$$1 \to \pi_1^{\acute{e}t}(\mathcal{C}_{\bar{T}_0}, 0_{\bar{T}_0}) \to \pi_1^{\acute{e}t}(\mathcal{C}, 0_{\bar{T}_0}) \to \pi_1^{\acute{e}t}(X, \bar{T}_0) \to 1, \tag{9}$$

which is a short exact sequence by [3], Corollaire X.2.2. Moreover, the section $\bar{s} : X \to \bar{\mathcal{C}}$ similarly gives rise to an action of $\pi_1^{\acute{e}t}(X, \bar{T}_0)$ on $\pi_1^{\acute{e}t}(\bar{\mathcal{C}}_{T_0}, \infty_{\bar{T}_0})^{\mathrm{ab}}$.

Step 3: We switch from $\bar{C}$ to its Jacobian. Define $\mathcal{J} \to X$ to be the abelian scheme representing the Picard functor of the scheme $\mathcal{C} \to X$ (see [4], Theorem 8.1). Note that $\mathcal{J}_T$ is the Jacobian of $\mathcal{C}_T$ for each $\mathbb{C}$-point T of X, and the generic fiber of $\mathcal{J}$ is J/L, the Jacobian of C/L. Let $f_\infty : \bar{C} \to J$ be the morphism (defined over L) given by sending each point $P \in \bar{C}(L)$ to the divisor class $[(P) - (\infty)]$ in $\mathrm{Pic}^0_L(\bar{C})$, which is identified with $J(L)$. By [4], Proposition 9.1, the induced homomorphism of étale fundamental groups $(f_\infty)_* : \pi_1^{\acute{e}t}(\bar{C}, \infty) \to \pi_1^{\acute{e}t}(J, 0)$ factors through an isomorphism $\pi_1^{\acute{e}t}(\bar{C}, \infty)^{\mathrm{ab}} \xrightarrow{\sim} \pi_1^{\acute{e}t}(J, 0)$. This induces an isomorphism $\pi_1^{\acute{e}t}(\bar{\mathcal{C}}_T, \infty_T)^{\mathrm{ab}} \xrightarrow{\sim} \pi_1^{\acute{e}t}(\mathcal{J}_T, 0_T)$ for each $T \in X$. Note that the composition of the section $\bar{s} : X \to \bar{\mathcal{C}}$ with f_∞ is the "zero section" $o : X \to \mathcal{J}$ mapping each T to the identity element $0_T \in \mathcal{J}_T$. Thus, the action of $\pi_1^{\acute{e}t}(X, \bar{T}_0)$ on $\pi_1^{\acute{e}t}(\mathcal{C}_{T_0}, \infty_{\bar{T}_0})^{\mathrm{ab}}$ coming from the splitting of (5) is the same as the action of $\pi_1^{\acute{e}t}(X, \bar{T}_0)$ on $\pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$ coming from the splitting of (9) induced by the section $o_* : \pi_1^{\acute{e}t}(X, \bar{T}_0) \to \pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{T}_0})$.

Step 4: We now show that this action on $\pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$ is isomorphic to a Galois action on $\pi_1^{\acute{e}t}(J_{\bar{L}}, 0)$ (and therefore on its ℓ-adic quotient $T_\ell(J)$). Let $\eta : \mathrm{Spec}(L) \to X$ denote the generic point of X. Note that we may identify $\pi_1^{\acute{e}t}(L, \bar{L})$ with G_L, and that $\eta_* : G_L \twoheadrightarrow \pi_1^{\acute{e}t}(X, \bar{\eta})$ is a surjection (in fact, it is the restriction homomorphism of Galois groups corresponding to the maximal algebraic extension of L unramified at all points of X). Also, the point $0 \in J_L$ may be viewed as a morphism $0 : \mathrm{Spec}(L) \to J_L$ which induces $0_* : G_L = \pi_1^{\acute{e}t}(L, \bar{L}) \to \pi_1^{\acute{e}t}(J_L, 0)$. Let $\bar{T}_0$ and $\bar{\eta}$ be geometric points over T_0 and η respectively. Then we have ([3], Corollaire X.1.4) an exact sequence of étale fundamental groups

$$\pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{\eta}}) \to \pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{\eta}}) \to \pi_1^{\acute{e}t}(X, \bar{\eta}) \to 1. \tag{10}$$

Changing the geometric basepoint of X from $\bar{T}_0$ to $\bar{\eta}$ (resp. changing the geometric basepoint of $\mathcal{J}$ from $0_{\bar{\eta}}$ to $0_{\bar{T}_0}$) non-canonically induces an isomorphism $\pi_1^{\acute{e}t}(X, \bar{\eta}) \xrightarrow{\sim} \pi_1^{\acute{e}t}(X, \bar{T}_0)$ (resp. an isomorphism $\pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{\eta}}) \xrightarrow{\sim} \pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{T}_0})$). Fix such an isomorphism $\varphi : \pi_1^{\acute{e}t}(X, \bar{\eta}) \xrightarrow{\sim} \pi_1^{\acute{e}t}(X, \bar{T}_0)$. Then we have the following commutative diagram, where all horizontal rows are exact:

$$\begin{array}{ccccccccc}
 & & & & & \overset{0_*}{\curvearrowleft} & & & \\
1 & \longrightarrow & \pi_1^{\acute{e}t}(J_{\bar{L}}, 0) & \longrightarrow & \pi_1^{\acute{e}t}(J_L, 0) & \longrightarrow & \pi_1^{\acute{e}t}(L, \bar{L}) & \longrightarrow & 1 \\
 & & \| & & \downarrow & & \downarrow{\scriptstyle \eta_*} & & \\
 & & \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{\eta}}, 0_{\bar{\eta}}) & \longrightarrow & \pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{\eta}}) & \overset{o_*}{\curvearrowleft} & \pi_1^{\acute{e}t}(X, \bar{\eta}) & \longrightarrow & 1 \\
 & & \downarrow{\scriptstyle \mathrm{sp}} & & \downarrow{\scriptstyle \wr} & & \downarrow{\scriptstyle \wr\, \varphi} & & \\
1 & \longrightarrow & \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0}) & \longrightarrow & \pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{T}_0}) & \overset{o_*}{\curvearrowleft} & \pi_1^{\acute{e}t}(X, \bar{T}_0) & \longrightarrow & 1
\end{array}$$

Here the vertical arrow from $\pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{\eta}})$ to $\pi_1^{\acute{e}t}(\mathcal{J}, 0_{\bar{T}_0})$ is a change-of-basepoint isomorphism chosen to make the lower right square commute, and $\mathrm{sp} : \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{\eta}}, 0_{\bar{\eta}}) \to \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$ is the surjective homomorphism induced by a diagram chase on the bottom two horizontal rows. Grothendieck's Specialization Theorem ([3], Corollaire X.3.9) states that sp is an isomorphism, which implies that the second row is also a short exact sequence. Thus, the action of $\pi_1^{\acute{e}t}(X, \bar{T}_0)$ on $\pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$ arising from the splitting of the lower row by o_* is isomorphic to the action of $\pi_1^{\acute{e}t}(X, \bar{\eta})$ on $\pi_1^{\acute{e}t}(\mathcal{J}_{\bar{\eta}}, 0_{\bar{\eta}})$ arising from the splitting of the middle row by o_*, via the isomorphism $\mathrm{sp} : \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{\eta}}, 0_{\bar{\eta}}) \to \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$. In turn, a simple diagram chase confirms that this action, after pre-composing with $\eta_* : \pi_1^{\acute{e}t}(L, \bar{L}) \twoheadrightarrow \pi_1^{\acute{e}t}(X, \bar{\eta})$, can be identified with the action of $\pi_1^{\acute{e}t}(L, \bar{L})$ on $\pi_1^{\acute{e}t}(J_{\bar{L}}, 0)$ arising from the splitting of the top row by 0_*. We denote this action by $\tilde{R} : G_L = \pi_1^{\acute{e}t}(L, \bar{L}) \to \mathrm{Aut}(\pi_1^{\acute{e}t}(J_{\bar{L}}, 0))$. Since the Tate module $T_\ell(J)$ may be identified with the maximal pro-ℓ quotient of $\pi_1^{\acute{e}t}(J_{\bar{L}}, 0)$, $\tilde{R}$ induces an action of G_L on $T_\ell(J)$, which we denote by $\tilde{R}_\ell : G_L \to \mathrm{Aut}(T_\ell(J))$. One can identify the symplectic pairing on $\pi_1(\mathcal{J}_{T_0}, 0_{T_0})$ with the Weil pairing on $T_\ell(J)$ via the results in [6], Chapter IV, §24. Therefore, the image of $\tilde{R}_\ell$ is a subgroup of $\mathrm{Sp}(T_\ell(J))$.

By the above construction, we may identify the maximal pro-ℓ quotient of $\pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$ with $H_1(\mathcal{C}_{T_0}, \mathbb{Z}) \otimes \mathbb{Z}_\ell$. Note that the isomorphism $\mathrm{sp} : \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{\eta}}, 0_{\bar{\eta}}) \xrightarrow{\sim} \pi_1^{\acute{e}t}(\mathcal{J}_{\bar{T}_0}, 0_{\bar{T}_0})$ induces an isomorphism of their maximal pro-ℓ quotients $\mathrm{sp}_\ell : T_\ell(J) \xrightarrow{\sim} H_1(\mathcal{C}_{T_0}, \mathbb{Z}) \otimes \mathbb{Z}_\ell$. By construction, the representation $\tilde{R}_\ell$ is isomorphic to the representation R_ℓ via sp_ℓ.

Step 5: It now suffices to show that $\tilde{R}_\ell = \rho_\ell$. To determine $\tilde{R}_\ell$, we are interested in the action of G_L on the group $\mathrm{Aut}_{J_{\bar{L}}}(Z)$ for each ℓ-power-degree covering $Z \to J_{\bar{L}}$. But each such covering is a subcovering of $[\ell^n] : J_{\bar{L}} \to J_{\bar{L}}$, so it suffices to determine the action of G_L on the group of translations $\{t_P | P \in J[\ell^n]\}$ for each n. Recall that $0_* : G_L \to \pi_1^{\acute{e}t}(J_L, 0)$ is induced by the inclusion of the L-point $0 \in J_L$. Thus, for any $\sigma \in G_L$, $0_*(\sigma)$ acts on any connected étale cover of J_L via σ acting on the coordinates of the points. Since $\tilde{R}(\sigma)$ is conjugation by $0_*(\sigma)$ on $\pi_1^{\acute{e}t}(J_{\bar{L}}, 0) \lhd \pi_1^{\acute{e}t}(J_L, 0)$, one sees that for each n, $0_*(\sigma)$ acts on $\{t_P | P \in J[\ell^n]\}$ by sending each t_P to $\sigma^{-1} t_P \sigma = t_{P^\sigma}$. Thus, G_L acts on the Galois group of the covering $[\ell^n] : J_{\bar{L}} \to J_{\bar{L}}$ via the usual Galois action on $J[\ell^n]$. This lifts to the usual action of G_L on $T_\ell(J)$, and we are done.

□

It is now easy to prove the main theorem.

Proof (of Theorem 1.1) . Recall that P_{2g+1} is the normal subgroup of $B_{2g+1} \cong \pi_1(X, T_0)$ corresponding to the cover $Y \to X$, and the function field of Y is $\mathbb{C}(\alpha_1, ..., \alpha_{2g+1}) = L_1$. It follows that the image of $\mathrm{Gal}(\bar{L}/L_1)$ under η_* is $\widehat{P}_{2g+1} \lhd \widehat{B}_{2g+1} \cong \pi_1^{\acute{e}t}(X, \bar{T}_0)$ (where $\widehat{P}_{2g+1}$ denotes the profinite completion of P_{2g+1}). Therefore, the statement of Theorem 2.1 with $\ell = 2$ implies that the image of $\mathrm{Gal}(\bar{L}/L_1)$ under R_2 is $\Gamma(2) \lhd \mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z}) \otimes \mathbb{Z}_\ell)$. It then follows from the statement of Lemma 2.2 that the image of $\mathrm{Gal}(\bar{L}/L_1)$ under ρ_2 is $\Gamma(2) \lhd \mathrm{Sp}(T_2(J))$.

□

3 Fields of 4-torsion

One application of Theorem 1.1 is that it allows us to obtain an explicit description of L_2. We will follow Yu's argument in [10].

Proposition 3.1. *We have*

$$L_2 = L_1(\{\sqrt{\alpha_i - \alpha_j}\}_{1 \le i < j \le 2g+1}).$$

Proof. For $n \ge 1$, let $\mathcal{B}_n$ denote the set of bases of the free $\mathbb{Z}/2^n\mathbb{Z}$-module $\mathcal{J}_{T_0}[2^n]$. Then it was shown in the proof of Theorem 1.1 that G_L acts on $\mathcal{B}_n$ through the map $R : \pi_1(X, T_0) \to \mathrm{Sp}(H_1(\mathcal{C}_{T_0}, \mathbb{Z})) = \mathrm{Sp}(H_1(\mathcal{J}_{T_0}, \mathbb{Z}))$ in the statement of Theorem 2.1, and the subgroup fixing all elements of $\mathcal{B}_n$ corresponds to $R^{-1}(\Gamma(2^n)) \lhd \pi_1(X, T_0)$. Hence, by covering space theory, there is a connected cover $X_n \to X$ corresponding to an orbit of $\mathcal{B}_n$ under the action of $\pi_1(X, T_0)$, and the function field of X_n is the extension of L fixed by the subgroup of G_L which fixes all bases of $J[2^n]$. Clearly, this extension is L_n.

Thus, the Galois cover $X_n \to X$ is an unramified morphism of connected affine schemes corresponding to the inclusion $L \hookrightarrow L_n$ of function fields.

Note that, setting $n = 1$, we get that X_1 is the Galois cover of X whose étale fundamental group can be identified with $R^{-1}(\Gamma(2)) \lhd \pi_1(X, T_0)$. Theorem 2.1 implies that $R^{-1}(\Gamma(2))$ is isomorphic to $\widehat{P}_{2g+1}$, the profinite completion of P_{2g+1}. For $n \geq 1$, the étale morphism $X_n \to X_1$ corresponds to the function field extension $L_n \supset L_1$, which by Corollary 1.2(c) has Galois group isomorphic to $\Gamma(2)/\Gamma(2^n)$. Therefore, X_n is the cover of X_1 whose étale fundamental group can be identified with a normal subgroup of $\widehat{P}_{2g+1}$ with quotient isomorphic to $\Gamma(2)/\Gamma(2^n)$.

In the proof of Corollary 2.2 of [8], it is shown that $\Gamma(2)/\Gamma(4) \cong (\mathbb{Z}/2\mathbb{Z})^{2g^2+g}$, and thus,

$$\mathrm{Gal}(L_2/L_1) \cong \Gamma(2)/\Gamma(4) \cong (\mathbb{Z}/2\mathbb{Z})^{2g^2+g}. \tag{11}$$

It is also clear from looking at a presentation of the pure braid group P_{2g+1} (see for instance [2], Lemma 1.8.2) that the abelianization of P_{2g+1} is a free abelian group of rank $2g^2 + g$. Therefore, its maximal abelian quotient of exponent 2 is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^{2g^2+g}$. Thus, $\widehat{P}_{2g+1}$ has a unique normal subgroup inducing a quotient isomorphic to $(\mathbb{Z}/2\mathbb{Z})^{2g^2+g}$. It follows that there is only one Galois cover of X_1 with Galois group isomorphic to $\Gamma(2)/\Gamma(4)$, namely X_2. The field extension $L_1(\{\sqrt{\alpha_i - \alpha_j}\}_{i<j}) \supset L_1$ is unramified away from the hyperplanes defined by $(\alpha_i - \alpha_j)$ with $i \neq j$ and is obtained from L_1 by adjoining $2g^2+g$ independent square roots of elements in $L_1^\times \backslash (L_1^\times)^2$. Therefore, $L_1(\{\sqrt{\alpha_i - \alpha_j}\}_{i<j})$ is the function field of a Galois cover of $X(2)$ with Galois group isomorphic to $(\mathbb{Z}/2\mathbb{Z})^{2g^2+g} \cong \Gamma(2)/\Gamma(4)$. It follows that this cover of X_1 is X_2, and that $L_1(\{\sqrt{\alpha_i - \alpha_j}\}_{i<j})$ is L_2, the function field of X_2.

□

4 Generalizations

As in Section 1, let k be an algebraic extension of $\mathbb{Q}$ which contains all 2-power roots of unity, and let K be the transcendental extension obtained by adjoining the coefficients of (1) to k. We will also fix the following notation. Let C_K be the hyperelliptic curve defined over K given by the equation (1), and let J_K be its Jacobian. For each $n \geq 0$, let K_n be the extension of K over which the 2^n-torsion of J_K is defined. Note that, analogous to the situation with C/L, the extension K_2 is $k(\alpha_1, ..., \alpha_{2g+1})$, which is Galois over K with Galois group isomorphic to S_{2g+1}. Let $\rho_{2,K} : \mathrm{Gal}(K_\infty/K) \to \mathrm{Sp}(T_2(J_K))$ be the homomorphism arising from the Galois action on the Tate module of J_K. We now investigate what happens to the Galois action when we descend from working over $\mathbb{C}$ to working over k. (In what follows, we canonically identify $T_2(J)$ with $T_2(J_K)$ and $\Gamma(2^n)$ with the level-2^n congruence subgroup of $\mathrm{Sp}(T_2(J_K))$ for each $n \geq 0$.)

Proposition 4.1. *The statements of Theorem 1.1, Corollary 1.2, and Proposition 3.1 are true when L and ρ_2 are replaced by K and $\rho_{2,K}$ respectively.*

Proof. For any $n \geq 0$, let $\theta_n : \mathrm{Gal}(L_\infty/L_n) \to \mathrm{Gal}(K_\infty/K_n)$ be the composition of the obvious inclusion $\mathrm{Gal}(L_\infty/L_n) \hookrightarrow \mathrm{Gal}(L_\infty/K_n)$ with the obvious restriction map $\mathrm{Gal}(L_\infty/K_n) \twoheadrightarrow \mathrm{Gal}(K_\infty/K_n)$. Let $\bar{\rho}_2^{(\infty)}$ (resp. $\bar{\rho}_{2,K}^{(\infty)}$) be the representation of $\mathrm{Gal}(L_\infty/L)$ (resp. $\mathrm{Gal}(K_\infty/K)$) induced from ρ_2 (resp. $\rho_{2,K}$) by the restriction homomorphism of the Galois groups. It is easy to check that $\bar{\rho}_2^{(\infty)} = \bar{\rho}_{2,K}^{(\infty)} \circ \theta_0$. It will suffice to show that θ_0 is an isomorphism.

First, note that for any $n \geq 0$, θ_n is injective by the linear disjointness of K_∞ and L_n over K_n. Now suppose that $n \geq 1$. Then, as in the proof of Corollary 1.2, the image under $\bar{\rho}$ of $\mathrm{Gal}(L_\infty/L_n)$ is the entire congruence subgroup $\Gamma(2^n)$. Therefore, since θ_n is injective, the image under $\bar{\rho}_K$ of $\mathrm{Gal}(K_\infty/K_n)$ contains $\Gamma(2^n)$. But since K contains all 2-power roots of unity, the Weil pairing is Galois invariant, and so the image of $\mathrm{Gal}(K_\infty/K_n)$ must also be contained in $\Gamma(2^n)$. Therefore, θ_n is an isomorphism for $n \geq 1$. Now, using Corollary 1.2(a) and the fact that $\mathrm{Gal}(K(\alpha_1, ..., \alpha_{2g+1})/K) \cong S_{2g+1}$, we get the commutative diagram below, whose top and bottom rows are short exact sequences.

$$\begin{array}{ccccccccc}
1 & \longrightarrow & \mathrm{Gal}(L_\infty/L_1) & \longrightarrow & \mathrm{Gal}(L_\infty/L) & \longrightarrow & S_{2g+1} & \longrightarrow & 1 \\
 & & \downarrow{\scriptstyle \theta_1} & & \downarrow{\scriptstyle \theta_0} & & \| & & \\
1 & \longrightarrow & \mathrm{Gal}(K_\infty/K_1) & \longrightarrow & \mathrm{Gal}(K_\infty/K) & \longrightarrow & S_{2g+1} & \longrightarrow & 1
\end{array}$$

By the Short Five Lemma, since θ_1 is an isomorphism, so is θ_0.

□

Remark 4.2. a) Suppose we drop the assumption that k contains all 2-power roots of unity. Then $\rho_{2,K}(G_K)$ is no longer contained in $\mathrm{Sp}(T_2(J))$ in general. However, the Galois equivariance of the Weil pairing forces the image of $\rho_{2,K}$ to be contained in the group of symplectic similitudes

$$\mathrm{GSp}(T_2(J)) := \{\sigma \in \mathrm{Aut}(T_2(J)) \mid E_2(P^\sigma, Q^\sigma) = \chi_2(\sigma) E_2(P, Q)\ \forall P, Q \in T_2(J)\},$$

where $E_2 : T_2(J) \times T_2(J) \to \lim_{\leftarrow n} \mu_{2^n} \cong \mathbb{Z}_2$ is the Weil pairing on the 2-adic Tate module of J, and $\chi_2 : G_K \to \mathbb{Z}_2^\times$ is the cyclotomic character on the absolute Galois group of K. Galois equivariance of the Weil pairing also implies that K_∞ contains all 2-power roots of unity. Thus, $K_\infty \supset K(\mu_{2^\infty})$, and the statements referred to in Proposition 4.1 still hold when we replace K with $K(\mu_{2^\infty})$.

Furthermore, if K contains $\sqrt{-1}$, the Weil pairing on $J[4]$ is Galois invariant, so the image of $\mathrm{Gal}(K_2/K_1)$ coincides with $\Gamma(2)/\Gamma(4) \lhd \mathrm{Sp}(J[4])$ and is therefore isomorphic to $\mathrm{Gal}(L_2/L_1)$. It follows that Proposition 3.1 still holds over $K(\sqrt{-1})$; that is,

$$K_2 = K_1(\sqrt{-1}, \{\sqrt{\alpha_i - \alpha_j}\}_{1 \leq i < j \leq 2g+1}). \tag{12}$$

b) In addition, suppose that k is finitely generated over $\mathbb{Q}$ (for example, a number field). We may specialize by assigning an element of k to each coefficient of the degree-$(2g+1)$ polynomial in (1), and defining the corresponding Jacobian

J_k/k and Galois representation $\rho_{2,k} : G_k \to \mathrm{Sp}(T_2(J_k))$. Then we may use Proposition 1.3 of [7] and its proof (see also [9]) to see that for infinitely many choices of $e_1, ..., e_{2g+1} \in k$, $\rho_{2,k}(G_k)$ can be identified with $\rho_{2,K}(G_K)$ from part (a). We have $\rho_{2,k}(\mathrm{Gal}(\bar{k}/k(\mu_{2^\infty}))) = \rho_{2,k}(G_k) \cap \mathrm{Sp}(T_2(J_k))$, and therefore, the statements referred to in Proposition 4.1 still hold over $k(\mu_{2^\infty})$. Similarly, Proposition 3.1 still holds over $k(\sqrt{-1})$.

Acknowledgements

I am grateful to Yuri Zarhin for his many ideas and suggestions. I would also like to thank the referee, who suggested that this material be presented in a separate paper, and whose corrections were helpful in improving the exposition.

References

[1] Norbert A'Campo. Tresses, monodromie et le groupe symplectique. *Commentarii Mathematici Helvetici*, 54(1):318–327, 1979.

[2] Joan S Birman. *Braids, links, and mapping class groups*, volume 82. Princeton University Press, 1974.

[3] Alexander Grothendieck et al. *Revêtements étales et groupe fondamental* (SGA 1), Lecture Notes in Mathematics, 224. Springer-Verlag, 1971.

[4] James S Milne. Jacobian varieties. In *Arithmetic geometry*, pages 167–212. Springer, 1986.

[5] David Mumford. Tata lectures on theta II: Jacobian theta functions and differential equations. *Progress in Mathematics*, 43, 1984.

[6] David Mumford. *Abelian varieties*, 2nd edition. Oxford University Press, 1974.

[7] Rutger Noot. Abelian varieties – Galois representations and properties of ordinary reduction. *Compositio Mathematica*, 97(1):161–172, 1995.

[8] Masatoshi Sato. The abelianization of the level d mapping class group. *Journal of Topology*, 3(4):847–882, 2010.

[9] J.-P. Serre. Lettres à Ken Ribet du 1/1/1981 et du 29/1/1981, Collected Papers, Vol. IV. Springer-Verlag, Berlin-Heidelberg, 1996, pp. 1-20.

[10] Jiu-Kang Yu. Toward a proof of the Cohen-Lenstra conjecture in the function field case. *preprint*, 1997.